

The ultraviolet properties of Luminous Infrared Galaxies at $z \sim 0.7$

Is there any evolution in their dust attenuation?

V. Buat¹, D. Marcillac², D. Burgarella¹, E. Le Floch^{h2,3*}, G. Rieke², T. T. Takeuchi⁴, J. Iglesias-Parámo⁵, and C. K. Xu⁶

¹ Observatoire Astronomique Marseille Provence, Laboratoire d'Astrophysique de Marseille, BP8, 133761 Marseille cedex 12, France

e-mail: veronique.buat@oamp.fr

² Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721

³ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

⁴ Astronomical Institute, Tohoku University, Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan

⁵ Instituto de Astrofísica de Andalucía, CSIC, 18008 Granada, SPAIN

⁶ California Institute of Technology, MC 405-47, 1200 East California Boulevard, Pasadena, CA 91125

ABSTRACT

Aims. The total infrared (TIR: 8-1000 μm) and far-ultraviolet (FUV: $\sim 1500\text{\AA}$) luminosity functions of galaxies and the related luminosity densities ρ_{TIR} and ρ_{FUV} are known to evolve with redshift from $z=0$ to $z \sim 1$ but with a different rate: the galaxy populations appear brighter in the past at both wavelengths but the evolution in TIR is larger than in FUV. It leads to an increase of the ratio of TIR to FUV luminosity densities $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ which can be interpreted as a global increase of the dust attenuation from $z=0$ to $z \sim 1$. Our aim is to understand the origin of this increase: is it entirely due to a variation of the dust attenuation with the luminosity of the galaxies as seen at $z=0$ or are properties of galaxies evolving with the redshift?

Methods. We focus on infrared galaxies more luminous than $L_{\text{TIR}} = 10^{11} L_{\odot}$ at $z \sim 0.7$ observed by SPITZER/MIPS and we measure their ultraviolet emission at 2310 $\text{\AA}$ from GALEX images. These Luminous InfraRed Galaxies (LIRGs) represent the bulk of the TIR luminosity density at intermediate redshift. The analysis of the ratio of TIR to FUV (rest-frame) luminosity ($L_{\text{TIR}}/L_{\text{FUV}}$) enables us to discuss their dust attenuation and to compare it to what is found in the nearby universe for galaxies of similar infrared luminosity and selected in the same way

Results. Some evolution of $L_{\text{TIR}}/L_{\text{FUV}}$ and therefore of dust attenuation is found: LIRGs at $z=0.7$ span a larger range of $L_{\text{TIR}}/L_{\text{FUV}}$ ratios than at $z=0$ and their mean dust attenuation at FUV wavelengths is found ~ 0.5 mag lower than for their local counterparts. The decrease of dust attenuation is found lower than that reported in other studies for bright galaxies selected in UV rest-frame at $z=1$ and 2. A semi-quantitative analysis is performed which accounts for the general increase of dust attenuation with the bolometric luminosity of galaxies: it is found that the slight decrease of dust attenuation for LIRGs at $z=0.7$ remains consistent with the increase of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ with redshift.

Key words. galaxies: evolution-dust: extinction-infrared: galaxies-ultraviolet: galaxies

1. Introduction

Rest-frame far-ultraviolet and thermal infrared surveys are commonly used to probe star formation in the universe and its evolution as a function of redshift z . The problem of dust attenuation of stellar light in galaxies is central in these analyses because FUV emission is directly affected by it and infrared emission takes its origin in this process. As a consequence the total infrared (TIR) to FUV flux ratio is a robust tracer of dust attenuation in star forming galaxies (e.g. Buat & Xu, 1996; Meurer et al., 1999; Gordon et al., 2000)

In a broad sense, both observations (FUV or TIR) have led to similar conclusions: a decrease of star formation rate (SFR) from $z=1$ to $z=0$ (Flores et al., 1999; Le Floch et al., 2005; Schiminovich et al., 2005) which is also found with other tracers of star formation (e.g. Hopkins & Beacom, 2006, and references therein). Each wavelength range is sensitive to a specific galaxy population and recovering all star formation from mono-wavelength observations (ultraviolet or infrared) appears to be difficult, even at low z . Buat et al. (2006) have shown that FUV and TIR surveys of the nearby universe do not exactly lead to the same sampling of galaxies.

This effect seems to be amplified at higher z . Thanks to recent surveys of SPITZER and GALEX the total infrared (TIR) and far-ultraviolet (FUV-1530Å) luminosity functions are now available from $z=0$ to $z=1$ (Arnouts et al., 2005; Le Floc'h et al., 2005); a strong evolution of both luminosity functions is seen but the ratio of luminosity densities ρ_{TIR}/ρ_{FUV} increases from $z=0$ to $z=1$ (Takeuchi et al., 2005b). This increase might be explained, at least qualitatively: dust attenuation is found to increase with the bolometric luminosity or SFR of galaxies in the nearby universe (Wang & Heckman, 1996; Buat & Burgarella, 1998; Sullivan et al., 2001; Hopkins et al., 2001; Martin et al., 2005; Buat et al., 2005) but also at higher z (Reddy et al., 2005; Le Floc'h et al., 2005; Bell et al., 2005). Therefore the general brightening of the galaxies when z increases, observed from the evolution of luminosity functions, may also induce an increase of the global dust attenuation.

Nevertheless, studies of galaxy samples at $z > 0$ have led to rather controversial results about the amount of dust obscuration in bright distant galaxies. Reddy et al. (2005) and Burgarella et al. (2006a) performed a FUV (rest-frame) like selection at $z=2$ and 1 respectively and found a dust attenuation about ten times lower than in the nearby universe. Conversely, samples selected in infrared by Choi et al. (2006) or Bell et al. (2005) led to attenuations consistent with the relations found at $z=0$ between the TIR to FUV flux ratio and the total SFR (or equivalently the luminosity of young stars). From GALEX and SWIRE observations of the ELAIS-N1 field Xu et al. (2006) found average trends consistent with no evolution between $z=0$ and $z=0.6$ of the TIR to FUV flux ratio of galaxies selected in FUV or at $24 \mu\text{m}$ except a decrease by a factor ~ 2 of the mean L_{TIR}/L_{FUV} for infrared selected galaxies with $L_{TIR} \sim 10^{11} L_{\odot}$. However, the detection rate in FUV (resp. $24 \mu\text{m}$) of galaxies selected at $24 \mu\text{m}$ (resp. FUV) was only 27 % (resp. 20%) and the analysis of Xu et al. (2006) almost entirely rely on stacking.

Here we analyse ultraviolet properties of Luminous InfraRed Galaxies (LIRGs with $L_{TIR} \geq 10^{11} L_{\odot}$) at medium z . Le Floc'h et al. (2005) have shown that LIRGs represent about half of the total infrared luminosity density at $z \sim 0.7$ and $\sim 70\%$ at $z \sim 1$ with only a minor contribution of Ultra Luminous InfraRed Galaxies (ULIRGs with $L_{TIR} \geq 10^{12} L_{\odot}$). On the contrary, most of UV selected galaxies are expected to have a lower TIR luminosity than LIRGs: at $z \sim 1$ Burgarella et al. (2006a) found that only 17% of their sample of Lyman Break Galaxies are detected by SPITZER at $24 \mu\text{m}$. A preliminary analysis of a FUV selection of galaxies at $z=0.7$ leads to only 10-15% of them with $L_{TIR} \geq 10^{11} L_{\odot}$ (Takeuchi et al., in preparation). However, even if they are not numerous, these UV selected galaxies also detected as LIRGs are found to be major contributors to the total star formation rate (Burgarella et al., 2006b).

We select our sample from the deep SPITZER/MIPS survey of the Chandra Deep Field South (Le Floc'h et al., 2005) focusing on redshift range 0.6-0.8. GALEX has also surveyed this area and very deep images are available. So we should have a very high detection rate of LIRGs with GALEX. In section 2, we present the data selection and the measurement of the ultraviolet emission of infrared sources. Then, in section 3, we analyse the ratio of the TIR to FUV luminosity (L_{TIR}/L_{FUV}) which is a tracer of dust attenuation in galaxies. A comparison between $z=0$ and $z=0.7$ is performed. In section 4 we compare our results to other studies. Section 5 is devoted to conclusions.

Throughout this article, we use the cosmological parameters $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$ and $\Omega_{\Lambda} = 0.7$. All magnitudes will be quoted in the AB system. The TIR luminosity L_{TIR} is defined on the wavelength range 8-1000 μm . The FUV luminosity L_{FUV} is defined as νL_{ν} with L_{ν} in $\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}$.

2. The data

2.1. Selection of the sample at $z \sim 0.7$

We start with the sample of galaxies in the Chandra Deep Field South (CDFS) used by Le Floc'h et al. (2005) to build TIR luminosity functions from $z=0$ to $z=1$. It consists of 2955 sources detected at 24 microns by MIPS with $f_{24} > 83 \mu\text{Jy}$ at the 80 % completeness limit (Papovich et al., 2004). As explained in Le Floc'h et al. (2005), several spectroscopic surveys (Le Fèvre et al., 2004; Vanzella et al., 2005; Szokoly et al., 2004) are primarily used to associate a spectroscopic redshift to the MIPS sources. Photometric redshifts from COMBO-17 (Wolf et al., 2004) are also used for sources at $z \leq 1.2$ and brighter than $R_{Vega} \sim 24$, where they are accurate enough ($\delta_z/(1+z) \leq 10\%$) to estimate quantities. We refer to Le Floc'h et al. (2005) for more details.

As discussed below, our strategy consists in measuring the NUV (2310Å) emission of these galaxies directly on the GALEX images at the position of the MIPS sources. We take a special care to avoid contaminations by sources in the close vicinity of MIPS ones which can also be ultraviolet emitters. At this aim, we apply strict selection criteria to the initial sample. All the steps followed to build the final sample and described in the sections 2 and 3 are summarized in Table 1. First, we work at $z \sim 0.7$ and only sources with a redshift comprised between 0.6 and 0.8 are kept; 623 galaxies are thus selected. A single optical counterpart must be found in COMBO-17 within 2 arcsec from the coordinates of the MIPS source. This tolerance radius was also adopted by Le Floc'h et al. (2005) for their identifications. This choice is motivated by the astrometrical precision of MIPS/ $24 \mu\text{m}$ images and the rather large FWHM of MIPS $24 \mu\text{m}$ PSF (~ 6 arcsec). It also accounts for a potential physical shift between the infrared and optical emission of disturbed objects. Here, we add an additional criterium: we exclude all MIPS sources associated to two or more optical sources within 2 arcsec. 65 sources are dropped and 558 sources are left.

We have not a complete sample of galaxies selected at $24 \mu\text{m}$ but we expect to have no strong bias in the selection. Our selection of "isolated" sources (i.e. with only one optical source within 2 arcsec) might avoid mergers (2 arcsec correspond to 14 kpc at $z=0.7$). Anyway, we cannot risk to have the FUV emission coming from another source than the MIPS one. Moreover, the reference sample at $z=0$ used for comparison is built in the same way (cf section 3.1). In order to check the effects of the exclusion of confused objects we have performed all the following analysis including confused sources both at high and low z and the results have been found unchanged.

Table 1. Number of sources at $0.6 < z < 0.8$ for the different selections applied to the original sample of galaxies selected at $24 \mu\text{m}$. Each row of the table corresponds to an additional selection applied to the original sample of 623 sources (from top to bottom). The final sample of LIRGs is described in the last row of the table

	number of sources at $24 \mu\text{m}$	NUV detections
original sample	623	–
a single optical counterpart	558	–
reliable NUV photometry	402	331
LIRGs	190	158

2.2. NUV measurements of the $24 \mu\text{m}$ sources

GALEX (Morrissey et al., 2005) observed the CDFS for 76 ks in both the FUV ($1530 \text{ \AA}$) and the NUV ($2310 \text{ \AA}$) as part of its deep imaging survey. The GALEX field of view (diameter 1.25 deg) is centered at $\alpha = 03^{\text{h}}32^{\text{m}}30.7^{\text{s}}$, $\delta = -27^{\circ}52'16.9''$. The resolution of the NUV image is measured and the FWHM of the PSF is found equal to 4.5 arcsec . Prior to any measurement we have applied a median filter 3×3 on the GALEX images which enlarges the PSF up to $\sim 6 \text{ arcsec}$ and makes it similar to that of the MIPS image.

We use DAOPHOT (Stetson, 1987) to measure the NUV emission at the location of the $24 \mu\text{m}$ sources. DAOPHOT was also used by Le Floc'h et al. (2005) to measure $24 \mu\text{m}$ fluxes. DAOPHOT is well suited for point sources (stellar fields). We must check that we can use it on our GALEX field. de Mello et al. (2004) have measured the size of UV selected galaxies at intermediate redshifts. In the redshift range $0.6-0.8 \sim 90\%$ of their observed galaxies have an effective (half light) optical radius R_e lower than 0.8 arcsec (which corresponds to $\sim 6 \text{ kpc}$ at $z=0.7$). If we assume a Gaussian distribution for the galaxy light, a galaxy with $R_e = 0.8 \text{ arcsec}$ convolved with the GALEX PSF (also assumed to be Gaussian) would appear with a FWHM $\sim 6.7 \text{ arcsec}$. The photometry is performed on the central 3 arcsec (i.e. 2 pixels for the GALEX images) which encloses 44% of the total energy for a pure PSF (6 arcsec after 3×3 median filtering, see above). For the largest objects ($R_e = 0.8 \text{ arcsec}$) the aperture of 3 arcsec would enclose 40% of the total energy. Therefore we estimate the photometric uncertainty assuming all the sources punctual to be at most 4%

The PSF was built using 10 stellar like objects with a FUV magnitude comprised between 17.5 and 19.8 mag . The GALEX field is rather dense with an average of ~ 0.06 galaxy per beam. The influence of close neighbours in the measure of NUV emission has been tested by adding artificial sources (ADDSTAR task): since we have excluded all the original MIPS sources with more than one optical source within 2 arcsec we simulated NUV point sources in the GALEX images located from 2 to 4 arcsec from the objects to be measured. As long as the galaxy has a magnitude lower or equal to $\text{NUV} = 24 \text{ mag}$ the contamination due to neighbours remains low (less than 0.4 mag in the worst cases) but when the source is fainter than $\text{NUV} = 25 \text{ mag}$ the contamination by neighbours can be very high (reaching 1 mag or more), whatever the magnitude and the location of the contaminating source is. Therefore we decided to exclude the sources whose measured NUV emission is fainter than $\text{NUV} = 24.5 \text{ mag}$ and which have at least one optical neighbour within 4 arcsec from the MIPS coordinates. At the end we are left with 402 sources for which the measure of NUV flux is considered as reliable. 156 galaxies are excluded because they are fainter than $\text{NUV} = 24.5 \text{ mag}$ and with an optical neighbour within 4 arcsec . As for the confused sources we have checked that including the sources with a non reliable NUV photometry (according to our strict criteria) does not modify the results of the subsequent analysis.

331 out of the 402 galaxies are detected in NUV. When the MIPS source is not detected by GALEX we put an upper limit on the NUV mag: $\text{NUV} = 26.2 \text{ mag}$. It corresponds to a completeness of 80% and a photometric error of $0.07 \pm 0.04 \text{ mag}$. This limit is obtained by simulations of 500 artificial sources added to the original GALEX image (ADDSTAR task). The NUV magnitudes are corrected for foreground Galactic extinction using the dust map of Schlegel et al. (1998) and the Galactic extinction curve from Cardelli et al. (1989).

It might be worth noting that only $\sim 20\%$ of the NUV sources at $z \sim 0.7$ have a MIPS counterpart brighter than $83 \mu\text{Jy}$. However, even if these galaxies are far from being a dominant population (in number) for a UV selection, their contribution to the total star formation rate of a UV selected sample is likely to be important because of their very high luminosity: Burgarella et al. (2006b) show that they account for $\sim 2/3$ of the total star formation rate of a sample of Lyman Break selected at $z \sim 1$. A full analysis of a FUV (rest-frame) galaxy sample at $z=0.7$ is underway (Takeuchi et al., in preparation).

3. The $L_{\text{TIR}}/L_{\text{FUV}}$ ratio of Luminous InfraRed Galaxies

We focus on Luminous InfraRed Galaxies (LIRGs, brighter than $10^{11} L_{\odot}$) because the SPITZER/MIPS sample is complete up to $z=0.8$ for $L_{\text{TIR}} > 10^{11} L_{\odot}$ and these LIRGs are found to be at the origin of the bulk of the TIR emission at medium z (Le Floc'h et al., 2005). Our aim is to compare the properties of a sample of LIRGs at $z=0.7$ to those of a reference sample taken at $z=0$ which will be defined in section 3.1. Indeed, from $z=0$ to $z=0.7$ ρ_{TIR} and ρ_{FUV} are found to increase by a factor equal to ~ 8 and ~ 4 respectively, leading to a net increase of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ by a factor ~ 2 (Takeuchi et al., 2005b). So, we can expect to find some evolution (if any) on the properties of galaxies in this redshift range.

We start by the description of the reference sample in order to define the quantities to be compared with the $z=0.7$ sample.

3.1. The reference sample at $z=0$

Buat et al. (2006) have built a sample of galaxies from the IRAS PSCz cross-correlated with the GALEX All Sky Imaging Survey (AIS) over more than 2000 deg². It is a flux limited sample ($f_{60} > 0.6$ Jy) of ~ 700 galaxies, most of them have a measured FUV flux at 1530 Å from the GALEX All sky Imaging Survey.

The total infrared (TIR) emission of the galaxies was estimated from their emission at 60 and 100 μm (see Buat et al. (2006) for more details). From this sample we select only LIRGs ($L_{\text{TIR}} > 10^{11} L_{\odot}$). We must restrict the sampled volume to be sure to detect all LIRGs within this volume. The limit $L_{\text{TIR}} = 10^{11} L_{\odot}$ corresponds to $L_{60} = 0.47 \cdot 10^{11} L_{\odot}$ for a mean L_{TIR}/L_{60} ratio of 2.13 (mean value found for the LIRGs of our reference sample). With a flux limit at 60 μm in the PSCz of 0.6 Jy, we must truncate the sample to $v < 16000 \text{ km s}^{-1}$. 98 LIRGs are selected, 91 have a measured FUV flux, 7 are not detected by GALEX in FUV and an upper limit at FUV = 20.5 mag is adopted corresponding to a 3σ detection limit for the GALEX-AIS survey (Morrissey et al., 2005). The criterium applied at $z=0$ to avoid confused sources was the absence of any neighbour within 1 arcmin from the IRAS source (Buat et al., 2006): at the distance of the selected LIRGs it corresponds to a projected distance larger than 14 kpc. Therefore we can assume that a similar criterium is applied at $z=0$ and $z=0.7$ (c.f. section 2.1) to select isolated sources.

3.2. The $z=0.7$ sample of LIRGS

We choose to work at $z \sim 0.7$ because there is an over density of galaxies at this redshift in the Chandra Deep Field South (e.g. Wolf et al., 2004). 190 galaxies of our sample of 402 galaxies are selected as Luminous InfraRed Galaxies with $L_{\text{TIR}} > 10^{11} L_{\odot}$, the mean redshift of the LIRGs sample is $\langle z \rangle = 0.70 \pm 0.05$. 158 out of the 190 LIRGs (i.e. 83%) are detected in NUV.

The total infrared (TIR) emission of these galaxies has been calculated by Le Floc'h et al. (2005) from their emission at 24 μm and we adopt the results of their calculations. At $z=0.7$ the GALEX NUV band (2310 Å) corresponds approximately to the GALEX FUV one (1530 Å) adopted for the reference sample (1358 Å for the rest-frame UV emission observed in NUV at $z=0.7$ against 1530 Å for the GALEX FUV band). The amplitude of K-corrections are investigated. The UV continuum is assumed to be well described by a power-law $f_{\lambda} \propto \lambda^{\beta}$ (e.g. Calzetti et al., 2000), where f_{λ} is expressed in $\text{erg cm}^{-2} \text{s}^{-1} \text{\AA}^{-1}$ (or $f_{\nu} \propto \nu^{-\beta-2}$ where f_{ν} is expressed in $\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$). $\beta = -1$ corresponds to a flat distribution in $\nu \times f_{\nu}$ (no K-correction). Unfortunately β cannot be measured for our sample at $z=0.7$. At $z=0$, β can be deduced from the FUV-NUV color measured by GALEX (e.g. Seibert et al., 2005). β is found to vary from -1.5 to 1 which induces $-0.03 \leq \log(\nu f_{\nu})_{\text{FUV rest-frame}} - \log(\nu f_{\nu})_{\text{NUV observed}} \leq 0.10$. Therefore we prefer not to apply any K-correction to the data: we consider the NUV fluxes observed at $z=0.7$ as rest-frame FUV fluxes directly comparable with the FUV fluxes observed at $z=0$.

3.3. $L_{\text{TIR}}/L_{\text{FUV}}$ distributions

$L_{\text{TIR}}/L_{\text{FUV}}$ is a robust tracer of dust attenuation in star forming galaxies. Quantitative estimates can be made regardless of details of the geometry and star formation history as long as galaxies are still forming stars actively (e.g. Buat & Xu, 1996; Meurer et al., 1999; Gordon et al., 2000; Calzetti et al., 2000). Buat et al. (2005) have performed calibrations for the GALEX bands. We reproduce their formulae for the FUV band:

$$A(\text{FUV}) [\text{mag}] = -0.0333 \left(\log \frac{L_{\text{TIR}}}{L_{\text{FUV}}} \right)^3 + 0.3522 \left(\log \frac{L_{\text{TIR}}}{L_{\text{FUV}}} \right)^2 + 1.1960 \left(\log \frac{L_{\text{TIR}}}{L_{\text{FUV}}} \right) + 0.4967 \quad (1)$$

$L_{\text{TIR}}/L_{\text{FUV}}$ has been found to increase with the bolometric luminosity of young stars in galaxies from $z=0$ to at least $z=2$ (e.g. Buat et al., 2006, and references in the introduction). As a matter of consequence, a comparison of $L_{\text{TIR}}/L_{\text{FUV}}$ distributions can only be valid for galaxies exhibiting the same luminosity distribution. Our selection of LIRGs at $z=0$ and at $z=0.7$ compensates for the evolution of the TIR luminosity function reported by Le Floc'h et al. (2005) which lead to brighter galaxies at higher z . Indeed the distributions of L_{TIR} in both samples are found very similar with $\langle \log(L_{\text{TIR}}/L_{\odot}) \rangle = 11.22 \pm 0.17$ at $z=0$ and $\langle \log(L_{\text{TIR}}/L_{\odot}) \rangle = 11.24 \pm 0.20$ at $z=0.7$. Therefore we can safely compare $L_{\text{TIR}}/L_{\text{FUV}}$ distributions from both samples.

The histograms of $L_{\text{TIR}}/L_{\text{FUV}}$ at $z=0$ and $z=0.7$ are presented in Fig. 1. We perform statistical tests accounting for non detections to compare the distributions (IRAF/stds/analysis/statistics/twosamp task). The two distributions are found to be drawn from different parent populations with a probability larger than 0.99. The Kaplan Meier estimates of their mean are slightly different: $\langle \log(L_{\text{TIR}}/L_{\text{FUV}})_{z=0.7} \rangle = 1.673 \pm 0.044$ and $\langle \log(L_{\text{TIR}}/L_{\text{FUV}})_{z=0} \rangle = 1.897 \pm 0.063$. Note that the uncertainty on K-corrections (cf. section 3.2) does not question the robustness of the result since it implies in most cases an under-estimate of L_{FUV} (at most 0.1 dex) and therefore an over-estimate of $L_{\text{TIR}}/L_{\text{FUV}}$ at $z=0.7$.

If we translate these mean values in quantitative measurements of dust attenuation at FUV wavelength with formula (1) we find $\langle A(\text{FUV}) \rangle = 3.33 \pm 0.08 \text{ mag}$ at $z=0.7$ and $\langle A(\text{FUV}) \rangle = 3.81 \pm 0.13 \text{ mag}$ at $z=0$. So a slight difference in the mean dust obscuration can be inferred from these data and the shapes of the distributions are significantly different: the distribution of $L_{\text{TIR}}/L_{\text{FUV}}$ at $z=0.7$ appears broader than at $z=0$ with an extension toward low $L_{\text{TIR}}/L_{\text{FUV}}$ values.

The difference found between the mean $L_{\text{TIR}}/L_{\text{FUV}}$ values is slight. Therefore one may question the presence of some systematic effects in the derivation of the total infrared luminosity from the observables. Indeed, at $z=0.7$ L_{TIR} is deduced from the flux observed at 24 μm (14 μm in the rest-frame at $z=0.7$) by Le Floc'h et al. (2005) using a set of local templates whereas at $z=0$ L_{TIR} is calculated with a combination of IRAS fluxes at 60 and 100 μm (Buat et al., 2006). Takeuchi et al. (2005a) have proposed a formula to derive L_{TIR} from the luminosity at 15 μm (rest-frame) fully consistent with the IRAS database. To check the robustness of our present results we used the calibration of Takeuchi et al. (2005a) to derive L_{TIR} at $z=0.7$ from the observed fluxes at 24 μm : we found




Fig. 1. L_{TIR}/L_{FUV} distribution for LIRGs at $z=0$ (black, hatched histogram) and $z=0.7$ (red, empty histogram), for both samples histograms of upper limits are plotted with dashed lines

the results unchanged with $\langle (L_{TIR}/L_{FUV})_{z=0.7} \rangle = 1.660 \pm 0.042$. One must remind, however, that our analysis is based on the hypothesis that local galaxy templates are valid to estimate the total infrared luminosity of high redshift systems (Le Flocc'h et al., 2005; Marcillac et al., 2006).

4. Discussion

4.1. Comparison with previous works

In Fig 2 is plotted the bolometric luminosity L_{bol} , defined as $L_{TIR} + L_{FUV}$, versus L_{TIR}/L_{FUV} for both samples at $z=0$ and $z=0.7$. The difference in the distributions is clearly seen with an extension to moderate L_{TIR}/L_{FUV} values at $z=0.7$ not present at $z=0$.

For comparison, we have also gathered the results of previous works on the variation of L_{TIR}/L_{FUV} at low and high z . In the nearby universe, we refer to Buat et al. (2006) and report here the mean relations found in their paper which are consistent with other studies of local galaxies (see Buat et al., 2006, for more discussions).

At higher z several recent studies are available. Choi et al. (2006) have selected galaxies at $z \sim 0.8$ at NIR+MIR wavelengths. By comparing the SFR deduced from the TIR emission and the strength of the emission lines they measured the extinction in the optical emission lines and found that the corresponding visual extinction varies as $A_V = 0.75 \log(L_{TIR}/L_{\odot}) - 6.35$ mag. To compare these results to the present work, we have to translate this visual extinction in the gaseous medium to an attenuation for the ultraviolet stellar continuum. To be consistent with Choi et al. (2006), we follow the Calzetti et al. recipe (Calzetti et al., 2000; Calzetti, 2001). For the emission lines we adopt a foreground like distribution with a Milky Way extinction curve for a diffuse medium (Cardelli et al., 1989):

$$A_V = 3.1 E(B - V)_g$$

where $E(B - V)_g$ is the color excess for the gas emission lines.

The color excess for the stellar continuum $E(B - V)_s$ is given by

$$E(B - V)_s = 0.44 E(B - V)_g$$

(Calzetti et al., 2000; Calzetti, 2001). Then using the reddening curve (Calzetti et al., 2000), it comes

$$A_{FUV} = k'(1530\text{\AA}) E(B - V)_s = 10.33 \times 0.44 \times (A_V/3.1)$$

with $k'(1530\text{\AA}) = 10.33$ which gives $A_{FUV} = 1.47 A_V$.

A_{FUV} is linked to L_{TIR}/L_{FUV} via formula (1). The result of these transformations is shown in Fig. 2, note that in the Choi et al. relation the x axis is L_{TIR} and not $L_{TIR} + L_{FUV}$: for the LIRG regime (the topic of the present work) the difference is very small.




Fig. 2. $L_{\text{TIR}}/L_{\text{FUV}}$ versus $L_{\text{bol}} = L_{\text{TIR}} + L_{\text{FUV}}$. Vertical right axis: the dust attenuation $A(\text{FUV})$ is calculated with formula (1) in the text. LIRGs are plotted with black 'plus' and arrows for our $z=0$ sample and red dots and arrows for our $z=0.7$ sample. Blue dashed line: FUV selected sample at $z=0$ from Buat et al. (2006), red solid line: FIR selected sample at $z=0$ from Buat et al. (2006). Dot dashed magenta line from Choi et al. (2006). The data from Reddy et al. (2005) are plotted with cyan lozanges, and those of Burgarella et al. (2006a,b) with green crosses

The relation of Choi et al. (2006) is consistent with the $z=0$ results at intermediate luminosity. For LIRG luminosities and higher the relation flattens and is below the relation found for FIR selected galaxies at $z=0$ (and consistent with the FUV selection at $z=0$). This trend is fully consistent with the present work and the presence of LIRGs exhibiting a lower $L_{\text{TIR}}/L_{\text{FUV}}$ ratio.

Therefore, a difference is found between the amount of dust attenuation in LIRGs selected in infrared surveys at $z=0$ and $z=0.7$. The mean value of $L_{\text{TIR}}/L_{\text{FUV}}$ varies by 0.2 dex which translates in a decrease of dust attenuation of ~ 0.5 mag from $z=0$ to $z=0.7$. The distribution of $L_{\text{TIR}}/L_{\text{FUV}}$ is broader at $z=0.7$ than at $z=0$: a population of LIRGs with a moderate $L_{\text{TIR}}/L_{\text{FUV}}$ ($\log(L_{\text{TIR}}/L_{\text{FUV}}) \sim 1 - 1.5$ corresponding to an attenuation of $\sim 2 - 3$ mag in FUV) appears at $z=0.7$, this population is not important at $z=0$.

A decrease of the dust attenuation can be linked to a lower metallicity in high redshift systems. Indeed, Liang et al. (2004) found a mean metallicity of LIRGs at $z > 0.4$ that is 0.3 dex lower as compared to that of local bright disks. The attenuation of the UV stellar continuum was found to be correlated with metallicity in starburst and normal galaxies at $z=0$ (Cortese et al., 2006; Heckman et al., 1998). Using the relation given by Cortese et al. (2006): $\log(L_{\text{TIR}}/L_{\text{FUV}}) = 1.37(12 + \log(O/H)) - 11.36$ we obtain a decrease of $L_{\text{TIR}}/L_{\text{FUV}}$ of 0.4 dex from $z=0$ to $z=0.7$ consistent with what is found in Fig. 2.

The observed decrease of dust attenuation in some LIRGs from $z=0$ to $z \sim 0.7$ might also be related to the evolution of morphological type for this galaxy population. From $z=0$ to $z > 0.5$ Melbourne et al. (2005) found a decrease of the number of peculiar/irregular systems exhibiting tidal features, assymetry or being obvious mergers as compared to spirals, this result is confirmed by Wang et al. (2006) and Bell et al. (2005). If disturbed galaxies are related to merging systems a larger dust attenuation is expected for them (e.g. Sanders & Mirabel, 1996) and their lower contribution to LIRGs at high z as compared to low z might imply a decrease of the mean dust attenuation for these galaxies.

At $z=2$ Reddy et al. (2005) studied star formation and dust obscuration of galaxies predominantly selected in optical (UV rest-frame). We select galaxies with $L_{\text{TIR}} > 10^{11} L_{\odot}$ and with $1.9 < z < 2.3$ from their sample (Reddy et al., 2006). In this redshift range the G-band corresponds to the FUV-band of GALEX in the galaxy rest-frame. We plot these selected data in Fig. 2: they appear to be distributed below our infrared selection. A comparison of the $L_{\text{TIR}}/L_{\text{FUV}}$ distributions is made difficult because of the obvious difference in the luminosity distributions of our samples at $z=0$ or 0.7 and the Reddy et al. sample at $z=2$ which contains a

relatively large number of ULIRGs ($L_{\text{TIR}} > 10^{12} L_{\odot}$). If the comparison is restricted to galaxies with $10^{11} < L_{\text{TIR}} < 10^{12} L_{\odot}$, the mean $\log(L_{\text{TIR}}/L_{\text{FUV}})$ found at $z=2$ is 0.9 corresponding to 1.85 mag, i.e. ~ 1.5 magnitudes lower than our mean value at $z=0.7$ for our selection of LIRGs at $24 \mu\text{m}$. The discrepancy seems to be lower for the brightest galaxies (ULIRGs). Given the low number of such bright galaxies in our sample at $z=0.7$ we cannot make any quantitative comparison; nevertheless in their study of the ELAIS-N1 field Xu et al. (2006) also found that dust obscuration in ULIRGs at $z=0.6$ is consistent with that obtained at $z=0$.

At $z=1$, Burgarella et al. (2006a) studied Lyman Break Galaxies (i.e. GALEX FUV dropouts) and also found a low obscuration for the galaxies they detected at $24 \mu\text{m}$ (only $\sim 15\%$ of their sample is detected at $24 \mu\text{m}$). In Fig. 2 we report their data (Burgarella et al., 2006b). The obscuration that they obtain is consistent with that found by Reddy et al. (2005) at $z=2$.

How to reconcile the results of Reddy et al. (2005) and Burgarella et al. (2006a) at $z \sim 1$ & $z \sim 2$ with ours at $z=0.7$? First we can invoke an evolution with the redshift but $z=1$ and $z=0.7$ are separated by only 1.5 Gyr and it is difficult to expect a large evolution during such a short timescale. The most natural explanation is to invoke selection effects since we have a selection at $24 \mu\text{m}$ ($15 \mu\text{m}$ rest-frame) whereas the selections of Reddy et al. (2005) and Burgarella et al. (2006a) are predominantly in the rest-frame ultraviolet. Nevertheless at $z=0$ Buat et al. (2006) find only a slight difference between a TIR and a FUV selection so we must assume a strong evolution of the properties of intrinsically luminous galaxies with z .

4.2. Do we expect a decrease of dust attenuation for LIRGs at $z \sim 0.7$?

At first glance the answer to this question is no: we do not expect a decrease of dust attenuation for bright galaxies at $z \sim 0.7$ since (as underlined in the introduction) the ratio of the luminosity densities $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ increases with z . Nevertheless, we observe a slight decrease of dust attenuation at a fixed L_{bol} and one must also account with the intrinsic brightening of the galaxies when z increases together with the increase of dust attenuation with the bolometric luminosity of galaxies observed at low and high z (cf. Fig. 2). Are all these trends consistent with each other?

We consider these issues according to a rather crude and semi-quantitative analysis (e.g. Xu et al., 2006). On one hand the evolution of luminosity densities in TIR and FUV from $z = 0$ to $z = 0.7$ has been quantified: Le Floc'h et al. (2005) found that ρ_{TIR} increases as $(1+z)^{3.9 \pm 0.4}$ and Schiminovich et al. (2005) obtained $(1+z)^{2.5 \pm 0.7}$ for the evolution of ρ_{FUV} . This gives an evolution of $\rho_{\text{TIR}}/\rho_{\text{FUV}} \propto (1+z)^{1.4 \pm 0.8}$. Therefore from $z = 0$ to $z = 0.7$ an increase of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ by a factor $2.1^{+1.1}_{-0.7}$ is obtained.

On the other hand, we can predict very crudely the evolution of dust attenuation of a typical L_{TIR}^* galaxy from the relation between dust attenuation and the bolometric luminosity of galaxies obtained at $z = 0$. This is regarded to represent the global trend of the evolution of galaxies selected in infrared, since Le Floc'h et al. (2005) showed that the evolution of TIR luminosity function is approximately described by the pure luminosity evolution (PLE) with a small density evolution. For the purpose of the discussion, let us perform a linear regression between $L_{\text{TIR}}/L_{\text{FUV}}$ and L_{bol} at $z = 0$: at this aim we take the mean values used to plot the solid and dashed lines at $z=0$ in Fig. 2 (see Buat et al. (2006) for more details) and we find

$$\log\left(\frac{L_{\text{TIR}}}{L_{\text{FUV}}}\right) = 0.64 \log(L_{\text{bol}}/L_{\odot}) - 5.5. \quad (2)$$

If we assume that L_{bol} is not very different of L_{TIR} (Buat et al., 2006), it translates to

$$\frac{L_{\text{TIR}}}{L_{\text{FUV}}} \propto L_{\text{TIR}}^{0.6}. \quad (3)$$

As mentioned above, Le Floc'h et al. (2005) found that the typical TIR luminosity L_{TIR}^* (the knee of the luminosity function) evolves as $(1+z)^{3.2^{+0.7}_{-0.2}}$, according roughly to the PLE. This gives

$$\frac{L_{\text{TIR}}}{L_{\text{FUV}}} \propto (1+z)^{1.9^{+0.4}_{-0.1}}. \quad (4)$$

Hence, if the $L_{\text{TIR}}/L_{\text{FUV}}-L_{\text{bol}}$ relation found at $z = 0$ is still valid at $z = 0.7$, a 'typical' TIR selected galaxy brightens with z and would have its $L_{\text{TIR}}/L_{\text{FUV}}$ increasing by a factor of ~ 3 (2.6–3.4). This factor is, however, slightly higher than that found for the evolution of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ (1.4–3.2).

These crude estimations show that a slight decrease of the mean $L_{\text{TIR}}/L_{\text{FUV}}$ from $z = 0$ to $z = 0.7$ is not inconsistent with the evolution of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ in the same redshift range. Moreover Xu et al. (2006) used a steeper regression between $L_{\text{TIR}}/L_{\text{FUV}}$ and L_{bol} (or equivalently SFR) and concluded to a larger discrepancy, the local relation between $L_{\text{TIR}}/L_{\text{FUV}}$ and L_{bol} predicting too much evolution of the cosmic dust attenuation.

Thus, we can conclude that the bulk of the variation of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ with z can be explained by *the increase of dust attenuation with the bolometric luminosity of galaxies and the brightening of the galaxies at high z* and that we do not need a global increase of dust attenuation in galaxies with a fixed luminosity. Moreover, the evolution of luminosity functions in TIR and FUV does not exclude a slight diminution of $L_{\text{TIR}}/L_{\text{FUV}}$ and hence of dust attenuation in individual galaxies from $z = 0$ to $z = 0.7$ ($\Delta(A_{\text{FUV}}) \simeq 0.5$ mag).

Note however that the much lower dust attenuation found in luminous UV selected galaxies at $z=1-2$ as compared to $z=0$ risks to be at odd with the evolution of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ with z as discussed above since the variation of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ does not seem to leave room for a strong decrease of dust attenuation in galaxies forming the bulk of the FUV and TIR luminosity densities. Nevertheless a substantial fraction of the galaxy samples of Reddy et al. (2005) and Burgarella et al. (2006a) are not detected in infrared and these non-detections must be accounted for a complete statistical analysis. A comparison at the same redshift of well controlled FUV and TIR (rest-frame) selected samples with a high detection rates at both wavelength (FUV and TIR) and for both samples will help to solve these issues. Such an analysis is in progress (Takeuchi et al. in preparation).

5. Conclusions

We have measured the FUV ($\sim 1500 \text{ \AA}$) rest-frame emission of a sample of 190 LIRGs at $z \sim 0.7$, 83% of these galaxies are detected in FUV. The ratio of the total infrared luminosity to the FUV one, $L_{\text{TIR}}/L_{\text{FUV}}$, is compared to that found for local galaxies of similar luminosity and also selected in far-infrared. The two samples at $z=0$ and $z=0.7$ are found to be drawn from different parent populations with a broader distribution of $L_{\text{TIR}}/L_{\text{FUV}}$ at $z=0.7$ and the presence of LIRGs with a moderate dust attenuation ($A(\text{FUV}) < 3 \text{ mag}$) as traced by $L_{\text{TIR}}/L_{\text{FUV}}$. A slight difference is found for $\langle L_{\text{TIR}}/L_{\text{FUV}} \rangle$ between the two samples (0.2 dex) implying a decrease of the mean dust attenuation of $\sim 0.5 \text{ mag}$ from $z=0$ to $z=0.7$. A lower dust attenuation in LIRGs at medium z might be related to the decrease of their metallicity. One may also invoke the increased fraction of spiral-like objects in the LIRG population from $z=0$ to $z > 0.5$ found by several authors. If we assume that disturbed objects are more affected by dust attenuation than undisturbed ones, dust attenuation is expected to decrease when z increases.

Nevertheless, the amplitude of the variation of the dust attenuation obtained for these infrared selected, bright galaxies is much lower than that reported for galaxies of similar luminosities at $z=1-2$, selected in UV-optical and detected in thermal infrared.

The intrinsic brightening of galaxies when redshift increases together with the well established variation of dust attenuation with the luminosity of galaxies explain the increase of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$ from $z=0$ to $z=0.7$. A slight diminution of dust attenuation in bright galaxies as observed here for the LIRG population remains consistent with the observed evolution of $\rho_{\text{TIR}}/\rho_{\text{FUV}}$.

Acknowledgements. V.B and D.B. gratefully acknowledge "Programme National Galaxie" and "Programme National Cosmologie" support for GALEX/SPITZER-MIPS science analysis. TTT has been supported by The 21st Century Center-of-Excellence Program "Exploring New Science by Bridging Particle-Matter Hierarchy", Tohoku University. Support for ELF's work was provided by NASA through the Spitzer Space Telescope Fellowship Program.

References

- Arnouts, S., Schiminovich, D., Ilbert, O., et al. 2005, *ApJ*, 619, L43
 Bell, E. F. 2003, *ApJ*, 586, 794
 Bell, E. F., Papovich, C., Wolf, C., et al. 2005, *ApJ*, 625, 23
 Buat, V., & Xu, C. 1996, *A&A*, 306, 61
 Buat, V., & Burgarella, D. 1998, *A&A*, 334, 772
 Buat, V., Iglesias-Páramo, J., Seibert, M., et al. 2005, *ApJ*, 619, L51
 Buat, V., et al. 2006 *ApJS*, in press (astro-ph/0609738)
 Burgarella, D., et al. 2006a, *A&A*, 450, 69
 Burgarella, D., Le Floc'h, E., Takeuchi, T.T. et al. 2006b, *MNRAS*, submitted
 Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, 533, 682
 Calzetti D. 2001, *PASP*, 113, 1449
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Choi, P.I., Yan, L., Im, M., et al. 2006, *ApJ*, 637, 227
 Cortese, L., Boselli, A., Buat, V., et al. 2006, *ApJ*, 637, 242
 Cowie, L.L., Songaila, A., Hu, E.M. & Cohen, J.G. 1996 *AJ* 112, 839
 de Mello, D.F., Wadadekar, Y., Dahlen, T., Casertano, S., & Gardner, J.P. 2006, *ApJ*, 131, 216
 Flores, H., et al. 1999, *ApJ*, 517, 148
 Gordon, K. D., Clayton, G. C., Witt, A. N., & Misselt, K. A. 2000, *ApJ*, 533, 236
 Heckman, T. M., Robert, C., Leitherer, C., Garnett, D. R., van der Rydt, F. 1998 *ApJ*, 503, 646
 Hopkins, A.M., Connolly, A.J., Haarsma, D.B., & Cram, L.E. 2001, *AJ* 122, 288
 Hopkins, A.M., Beacom, J.F. 2006, *astro-ph/0601463*
 Le Fèvre, O., Vettolani, G., Paltani, S., et al. 2004, *A&A*, 428, 1043
 Le Floc'h, E., Papovich, C., Dole, H., et al. 2005, *ApJ*, 632, 169
 Liang, Y. C., Hammer, F., Flores, et al. 2004, *A&A* 423, 867
 Marillac, D., Elbaz, D., Charlot, S. et al. 2006, *A&A* 458, 369
 Martin, D. C., Seibert, M., Buat, V., et al. 2005, *ApJ*, 619, L59
 Melbourne, J., Koo, D. C., & Le Floc'h, E. 2005, *ApJ*, 632, L65
 Meurer, G.R., Heckman, T.M., Calzetti, D. 1999, *ApJ*, 521, 64
 Morrissey, P., Schiminovich, D., Barlow, T.A., et al. 2005, *ApJ*, 619, L7
 Pérez-González, P. G., Rieke, G.H., Egami, E., et al. 2005, *ApJ*, 630, 82
 Reddy, N. A., Erb, D. K., Steidel, C. C., et al. 2005, *ApJ*, 633, 748
 Reddy, N. A., Steidel, C. C., Erb, et al. 2006, *astro-ph/0609296* Saunders, W., et al. 2000, *MNRAS*, 317, 55
 Sanders, D.B., Mirabel, I.F. 1996, *ARA&A*, 34, 749
 Schiminovich, D., Ilbert, O., Arnouts, S., et al. 2005, *ApJ*, 619, L47
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Seibert, M., Martin, D.C., Heckman, T.M., et al. 2005, *ApJ*, 619, L55
 Stetson, P.B. 1987, *PASP*, 99, 191
 Sullivan, M., Mobasher, B., Chan, et al. 2001, *ApJ* 558, 72
 Szokoly, G.P., Bergeron, J., Hasinger, G., et al. 2004, *ApJS*, 155, 271
 Takeuchi, T. T., Buat, V., Iglesias-Páramo, J., Boselli, A. & Burgarella, D. 2005a, *A&A*, 440, L17
 Takeuchi, T. T., Buat, V., & Burgarella, D. 2005b, *A&A*, 440, L17
 Vanzella, E., Cristiani, S., Dickinson, M., et al. 2005, *A&A*, 434, 53
 Wang, B., & Heckman, T. M. 1996, *ApJ*, 457, 645
 Wang, J.L., Xia, X.Y., Mao, S., et al. 2006, *ApJ*, 649, 722
 Wolf, C., Meisenheimer, K., Kleinheinrich, M., et al. 2004, *A&A*, 421, 913
 Xu, C. K., Shupe, D., Buat, V., et al. 2006, *ApJS*, submitted